arXiv:1508.06458v1 [math.AT] 26 Aug 2015

# NONEXISTENCE OF ALMOST COMPLEX STRUCTURES ON THE PRODUCT $S^{2m} \times M$

PRATEEP CHAKRABORTY AND AJAY SINGH THAKUR

Abstract. In this note we give a necessary condition for having an almost complex structure on the product $S^{2m} \times M$, where $M$ is a connected orientable closed manifold. We show that if the Euler characteristic $\chi(M) \neq 0$, then except for finitely many values of $m$, we do not have almost complex structure on $S^{2m} \times M$. In the particular case when $M = \mathbb{CP}^n, n \neq 1$, we show that if $n \not\equiv 3 \pmod 4$ then $S^{2m} \times \mathbb{CP}^n$ has an almost complex structure if and only if $m = 1, 3$. As an application we obtain conditions on the nonexistence of almost complex structure on Dold manifolds.

## 1. Introduction

Recall that an oriented manifold $X$ has an almost complex structure (a.c.s. for short) if there is a complex vector bundle $\xi$ such that the underlying real bundle $\xi_{\mathbb{R}}$ of $\xi$ is isomorphic to the tangent bundle $\tau_X$ of $X$ as oriented bundles.

It is well known [1] that the only even dimensional spheres admitting an a.c.s are $S^2$ and $S^6$. The product of even dimensional spheres, $S^{2m} \times S^{2n}$ with $m, n \neq 0$, has an a.c.s. if and only if $(m, n) = (1, 1), (1, 3), (3, 1), (1, 2), (2, 1)$ or $(3, 3)$ (see, for example, [2] and [9]). In [11], Tang showed that $S^{2m} \times \mathbb{CP}^2$ has an a.c.s. if and only if $m = 1, 3$, and $S^{2m} \times \mathbb{CP}^3$ has an a.c.s. if and only if $m = 1, 2, 3$.

In this note we deal with the question of existence of an a.c.s. on the general product of the form $S^{2m} \times M$, where $M$ is a connected orientable closed manifold of even dimension and $m \neq 0$. We have the following main result.

**Theorem 1.1.** *If $S^{2m} \times M$ has an a.c.s. then $2^r \cdot (m-1)!$ divides the Euler characteristic $\chi(S^{2m} \times M)$, where $2^r$ is the highest power of 2 dividing $m$.*

In the special case when $M = \mathbb{CP}^n$, we have the following result.

**Theorem 1.2.** *Let $n > 1$ and $n \not\equiv 3 \pmod 4$. Then $S^{2m} \times \mathbb{CP}^n$ has a.c.s. if and only if $m = 1, 3$.*

We also observe that $S^2 \times \mathbb{CP}^1$ has exactly two almost complex structures whereas $S^2 \times \mathbb{CP}^2$ and $S^4 \times \mathbb{CP}^3$ have infinitely many almost complex structures. As an application we obtain a result on the nonexistence of a.c.s. on Dold manifolds which strengthens the result obtained in [12].

For any real or complex vector bundle $\xi$ over a $CW$-complex $X$, we shall denote its stable class in $\widetilde{KO}(X)$ or $\widetilde{K}(X)$ by $[\xi]$. Let $Vect_n(X)$ be the set of all isomorphism classes of $n$-rank complex vector bundles over $X$. One of the stability properties of vector bundles over $X$ is that if $X$ is of dimension $2n$ then the map $Vect_n(X) \to \widetilde{K}(X)$ which takes the complex vector bundle $\xi$ to its stable class $[\xi]$ is bijective (see [7, Chapter 9, Theorem 1.5]). If $\rho : \widetilde{K}(X) \to \widetilde{KO}(X)$ is the realization map then the proof of our result is based on the following theorem.

---

2010 *Mathematics Subject Classification.* 57R15 (57R20).

*Key words and phrases.* Almost complex structure, Characteristic classes, Dold manifold.

The research of second author is supported by DST-Inspire Faculty Scheme(IFA-13-MA-26).

**Theorem 1.3.** ([9, Theorem 1.1] and [14, Lemma 1,1]) *A $2n$-dimensional connected oriented manifold $X$ admits an a.c.s. if and only if there exists an element $a \in \widetilde{K}(X)$ such that $\rho(a) = [\tau_X]$ and $n$th-Chern class $c_n(a)$ equals to the Euler class $e(X)$.* □

In this note, if $X$ is a complex manifold then the holomorphic tangent bundle of $X$ will be denoted by $T_X$. All the manifolds $M$ we consider here will be connected, closed, orientable and of dimension $2n$.

## 2. Almost complex structures on $S^{2m} \times M$

We first fix our notations. Let $\nu$ be the canonical complex line bundle over $S^2$. Let $g = (\nu - 1) \in \widetilde{K}(S^2)$ be a generator. Then by the Bott-Periodicity, $g^m \in \widetilde{K}(S^{2m})$ is a generator. We fix the generator $y_m \in H^{2m}(S^{2m};\mathbb{Z})$ such that the total Chern class $c(g^m) = 1 + (m-1)! \cdot y_m$. The Bott Integrability Theorem [7, Chapter 20, Corollary 9.8] says that for any $a \in \widetilde{K}(S^{2m})$, the top Chern class $c_m(a)$ is divisible by $(m-1)!$. (Here divisibility is in the sense that $c_m(a) = (m-1)! \cdot e_i$ for some $e_i \in H^{2m}(S^{2m};\mathbb{Z})$).

In the following proposition we shall give a condition on the Chern classes of any vector bundle over the smash product $S^{2m} \wedge M$.

**Proposition 2.1.** *For an element $a \in \widetilde{K}(S^{2m} \wedge M)$, each Chern class $c_i(a)$ is divisible by $(m-1)!$.*

*Proof.* Let $a \in \widetilde{K}(S^{2m} \wedge M)$. Then by the Bott-Periodicity, $a = g^m \otimes (\beta - n)$ for some complex vector bundle $\beta$ of rank $n$ over $M$. Let $c(\beta) = \prod_k (1 + s_k)$ be the formal factorization obtained by using the splitting principle. As in the proof of Lemma 2.1(2) [10], we can formally write the total Chern class of $a$ as follow

$$\begin{aligned} c(a) = {} & c(g^m \otimes (\beta - n)) \\ = {} & \textstyle\prod_k (1 + (m-1)! \cdot y_m((1+s_k)^{-m} - 1)) \qquad (1) \\ = {} & 1 + (m-1)! \cdot y_m(\textstyle\sum_{i \geq 1} (-1)^i \binom{m+i-1}{i} (\sum_k s_k^i)). \end{aligned}$$

The last equality in Equation 1 is due to the fact that $y_m^2 = 0$. The summation $\sum_k s_k^i$ can be expressed as a polynomial in Chern classes of $\beta$ with integer coefficients (by Newton-Girard formula). From here it is clear that each Chern class $c_i(a)$ is divisible by $(m-1)!$. □

Next we shall obtain conditions on Chern classes of any element in $\widetilde{K}(S^{2m} \times M)$. Consider the following split exact sequence

$$0 \to \widetilde{K}(S^{2m} \wedge M) \to \widetilde{K}(S^{2m} \times M) \to \widetilde{K}(S^{2m}) \oplus \widetilde{K}(M) \to 0.$$

Hence any element $a \in \widetilde{K}(S^{2m} \times M)$ can be written as $a = a_1 + a_2 + a_3$, where $a_1 \in \widetilde{K}(S^{2m} \wedge M)$, $a_2 \in \widetilde{K}(S^{2m})$ and $a_3 \in \widetilde{K}(M)$. In the following proposition we choose an orientation on $S^{2m} \times M$.

**Proposition 2.2.** *For any $a \in \widetilde{K}(S^{2m} \times M)$, all the Chern numbers of $a$ are divisible by $(m-1)!$.*

*Proof.* Let $a = a_1 + a_2 + a_3$ be as described above. Let the total Chern class $c(a_1) = 1 + y_m x_1 + y_m x_2 + \cdots + y_m x_n$, $c(a_2) = 1 + b y_m$ and $c(a_3) = 1 + z_1 + z_2 + \cdots + z_n$ where $x_i, z_i \in H^{2i}(M;\mathbb{Z})$ and $b$ is an integer. Since $a_1 \in \widetilde{K}(S^{2m} \wedge M)$, by Proposition 2.1, it is clear that each $y_m x_i$ is divisible by $(m-1)!$. Also since $a_2 \in \widetilde{K}(S^{2m})$, by Bott Integrability Theorem, we have that $b$ is divisible by $(m-1)!$. Now the proof of the proposition follows easily. □

To prove our main result consider the following commutative diagram

$$\begin{tikzcd}
\widetilde{K}(S^{2m} \times M) \arrow[d, "\rho"] & = & \widetilde{K}(S^{2m} \wedge M) \arrow[d, "\rho"] & \oplus & \widetilde{K}(S^{2m}) \arrow[d, "\rho"] & \oplus & \widetilde{K}(M) \arrow[d, "\rho"] \\
\widetilde{KO}(S^{2m} \times M) & = & \widetilde{KO}(S^{2m} \wedge M) & \oplus & \widetilde{KO}(S^{2m}) & \oplus & \widetilde{KO}(M)
\end{tikzcd}$$

In view of Theorem 1.3, if $a \in \widetilde{K}(S^{2m} \times M)$ gives an a.c.s. on $S^{2m} \times M$ then $\rho(a) = [\tau_{S^{2m} \times M}]$ and the top Chern class $c_{m+n}(a) = e(S^{2m} \times M)$. If we write $a = a_1 + a_2 + a_3$ then using the fact that $[\tau_{S^{2m} \times M}] = [\tau_M]$, we have $\rho(a_1) = 0$, $\rho(a_2) = 0$, $\rho(a_3) = [\tau_M]$.

**Proof of Theorem 1.1.** Suppose $a \in \widetilde{K}(S^{2m} \times M)$ gives an a.c.s. on $S^{2m} \times M$. From Proposition 2.2 and as $c_{m+n}(a) = e(S^{2m} \times M)$, it follows that $(m-1)!$ divides $\chi(S^{2m} \times M)$. Hence in the case when $m$ is odd, the proof of the theorem is complete.

Next assume that $m = 2p$. As above we write $a = a_1 + a_2 + a_3$. First observe that the realization map $\rho : \widetilde{K}(S^{4p}) \to \widetilde{KO}(S^{4p})$ is injective, and since $\rho(a_2) = 0$, we have $a_2 = 0$. Next note that $\rho(a_1) = 0$ and this implies that $a_1 + \overline{a_1} = 0$. Hence $2c_{2i}(a_1) = 0$ for $i > 0$ as the non-trivial cup-products in the cohomology of the smash product $S^{4p} \wedge M$ are zero. Therefore the top Chern class

$$c_{2p+n}(a) = \sum_{2i+1+j=2p+n} c_{2i+1}(a_1)c_j(a_3). \tag{2}$$

Now consider the element $a_1' \in \widetilde{K}(S^{4p+2} \wedge M)$ given as $a_1' = g \otimes a_1$. We can write $a_1 = g^{2p} \otimes (\beta - n)$ for some complex vector bundle $\beta$ of rank $n$ over $M$ and let $c(a_1) = 1 + y_{2p}x_1 + y_{2p}x_2 + \cdots + y_{2p}x_n$. Then as in the proof of Proposition 2.1, the total Chern class of $a_1'$ will be

$$c(a_1') = c(g \otimes a_1) = 1 + y_{2p+1}(\sum_{i \geq 1}(2p+i)x_i).$$

Since $a_1' \in \widetilde{K}(S^{4p+2} \wedge M)$, by Proposition 2.1, we have that each $(2p+i)y_{2p+1}x_i$ is divisible by $(2p)!$. This immediately implies that $(2p)!$ divides $(2p+i)y_{2p}x_i$. Hence for each odd $i$, $(2p)!$ will divide

$$(\prod_j (2p+j)) \cdot y_{2p}x_i,$$

where the product varies over odd $j$ and $1 \leq j \leq n$. Hence by Equation 2 we have that $(2p)!$ divides

$$(\prod_j (2p+j)) \cdot \chi(S^{4p} \times M),$$

where $j$ is odd and $1 \leq j \leq n$. Now the proof of the theorem follows from the fact that $(2p-1)!$ divides $\chi(S^{4p} \times M)$. □

From Theorem 1.1 we can observe that for a given $M$ with $\chi(M) \neq 0$, except for finitely many values of $m$, we do not have a.c.s on $S^{2m} \times M$. In particular we have the following corollary whose proof follows immediately.

**Corollary 2.3.** *Let* $\chi(M) \not\equiv 0 \pmod 4$ *or* $\chi(M)$ *be a power of two. Then* $S^{2m} \times M$ *does not have an a.c.s. for* $m \neq 1, 2, 3$. □

The restriction on $m$ in Corollary 2.3 is the best possible as we know that $S^2 \times \mathbb{CP}^n$, $S^6 \times \mathbb{CP}^n$, $S^4 \times S^2$ and $S^4 \times \mathbb{CP}^3$ (see Example 3.10 below) admits an a.c.s.

## 3. Almost complex structure on $S^{2m} \times \mathbb{CP}^n$

In this section we shall deal with the case when $M$ is a complex projective space $\mathbb{CP}^n$. By Corollary 2.3 we know that in the case when $n \not\equiv 3 \pmod 4$ or $n+1 = 2^k$ for some $k \geq 0$, the product $S^{2m} \times \mathbb{CP}^n$ does not admit a.c.s. for $m \neq 1, 2$ and 3. We shall prove that if $n \not\equiv 3 \pmod 4$ and $n > 1$ then $S^4 \times \mathbb{CP}^n$ also does not admit a.c.s.

Let $H$ be the canonical line bundle over $\mathbb{CP}^n$. Let $x = c_1(H) \in H^2(\mathbb{CP}^n; \mathbb{Z})$ be a generator. Let $\eta = H - 1 \in \widetilde{K}(\mathbb{CP}^n)$ and $r = [n/2]$. For $\alpha \in \widetilde{K}(S^{2m})$ and $\beta \in \widetilde{K}(\mathbb{CP}^n)$ we shall write the external product $\alpha \otimes \beta \in \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n)$ as $\alpha\beta$. In view of Lemma 3.5 [4] we have the following lemma whose proof follows from the Bott-Periodicity.

**Lemma 3.1.** *Each of the following system of elements form an integral basis of* $\widetilde{K}(S^{2m} \wedge \mathbb{CP}^n)$.

*(i)* $g^m\eta$, $g^m\eta(\eta+\bar\eta)$, $\cdots$, $g^m\eta(\eta+\bar\eta)^{r-1}$, $g^m(\eta+\bar\eta)$, $g^m(\eta+\bar\eta)^2$, $\cdots$, $g^m(\eta+\bar\eta)^r$, *and also in case* $n$ *is odd,* $g^m\eta^{2r+1} = g^m\eta(\eta+\bar\eta)^r$;

*(ii)* $g^m\eta$, $g^m\eta(\eta+\bar\eta)$, $\cdots$, $g^m\eta(\eta+\bar\eta)^{r-1}$, $g^m(\eta-\bar\eta)$, $g^m(\eta-\bar\eta)(\eta+\bar\eta)$, $\cdots$, $g^m(\eta-\bar\eta)(\eta+\bar\eta)^{r-1}$, *and also in case* $n$ *is odd,* $g^m\eta^{2r+1} = g^m\eta(\eta+\bar\eta)^r$. □

Let $w_k = g^m(H^k - 1) - \overline{g^m(H^k-1)} \in \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n)$. Similar to Proposition 4.3 [13], we have the following three propositions.

**Proposition 3.2.** *Let* $m \equiv 1, 3 \pmod 4$. *Then the kernel of the realization map* $\rho : \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n) \to \widetilde{KO}(S^{2m} \wedge \mathbb{CP}^n)$ *is freely generated by* $w_1, w_2, \cdots, w_r$.

*Proof.* It is clear that all $w_k$'s are in the kernel of the realization map. Further it can be proved using the the fact that $\bar g = -g$ in $\widetilde{K}(S^2)$ and using induction on $k$ that

$$g^m(\eta+\bar\eta)^k = w_k + (\text{linear combination of } w_1, \cdots, w_{k-1}).$$

Now using the structure of $\widetilde{KO}^{-2m}(\mathbb{CP}^n)$ (see [5]) and the fact that the $\mathbb{Q}$-linear map

$$\rho \otimes Id : \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n) \otimes_{\mathbb{Z}} \mathbb{Q} \to \widetilde{KO}(S^{2m} \wedge \mathbb{CP}^n) \otimes_{\mathbb{Z}} \mathbb{Q}$$

is surjective, we can see that $w_1, \cdots, w_k$ freely generates the kernel of $\rho$. □

For next two propositions we first note that if $m$ is even then

$$g^m(\eta-\bar\eta)(\eta+\bar\eta)^{k-1} = w_k + (\text{linear combination of } w_1, \cdots, w_{k-1}),$$

whose proof can again be given using induction on $k$. Next consider the following commutation diagram,

$$\begin{tikzcd}
\widetilde{K}(S^{2m} \wedge S^{2n}) \arrow[r] \arrow[d, "\rho"] & \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n) \arrow[r] \arrow[d, "\rho"] & \widetilde{K}(S^{2m} \wedge \mathbb{CP}^{n-1}) \arrow[d, "\rho"] \\
\widetilde{KO}(S^{2m} \wedge S^{2n}) \arrow[r] & \widetilde{KO}(S^{2m} \wedge \mathbb{CP}^n) \arrow[r] & \widetilde{KO}(S^{2m} \wedge \mathbb{CP}^{n-1})
\end{tikzcd}$$

Now the proof of the following two propositions follow from the fact that the realization map $\rho : \widetilde{K}(S^{2l}) \to \widetilde{KO}(S^{2l})$ is nonzero for $l \not\equiv 3 \pmod 4$.

**Proposition 3.3.** *Let* $m \equiv 0 \pmod 4$. *Then the kernel of the realization map* $\rho : \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n) \to \widetilde{KO}(S^{2m} \wedge \mathbb{CP}^n)$ *is freely generated by*

(1) $w_1, w_2, \cdots, w_r$ *when* $n$ *is even;*
(2) $w_1, w_2, \cdots, w_r, g^m\eta^n$, *when* $n \equiv 3 \pmod 4$;

(3) $w_1, w_2, \cdots w_r, 2g^m\eta^n$, *when* $n \equiv 1 \pmod 4$. ☐

**Proposition 3.4.** *Let* $m \equiv 2 \pmod 4$. *Then the kernel of the realization map* $\rho : \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n) \to \widetilde{KO}(S^{2m} \wedge \mathbb{CP}^n)$ *is freely generated by*

(1) $w_1, w_2, \cdots, w_r$ *when $n$ is even;*
(2) $w_1, w_2, \cdots, w_r, g^m\eta^n$, *when* $n \equiv 1 \pmod 4$*;*
(3) $w_1, w_2, \cdots w_r, 2g^m\eta^n$, *when* $n \equiv 3 \pmod 4$. ☐

Next we shall describe the Chern classes of elements in the Kernel of the realization map $\rho : \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n) \to \widetilde{KO}(S^{2m} \wedge \mathbb{CP}^n)$. Using Lemma 2.1(2) [10] and by the fact that $y_m^2 = 0$, we can easily compute the total Chern class of $w_k$. When $m$ is even then

$$c(w_k) = 1 - (m-1)! \cdot \sum_{i \geq 1} 2\binom{m+2i-2}{2i-1} k^{2i-1} y_m x^{2i-1}.$$

When $m$ is odd we have

$$c(w_k) = 1 + (m-1)! \cdot \sum_{i \geq 1} 2\binom{m+2i-1}{2i} k^{2i} y_m x^{2i}.$$

To compute the Chern class of $g^m\eta^n \in \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n)$, consider the following exact sequence

$$0 \to \widetilde{K}(S^{2m} \wedge S^{2n}) \to \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n) \to \widetilde{K}(S^{2m} \wedge \mathbb{CP}^{n-1}).$$

The element $g^m\eta^n \in \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n)$ is a generator of the kernel of the last map in the above sequence. Hence the total Chern class of $g^m\eta^n$ is as follows:

$$\begin{aligned} c(g^m\eta^n) &= 1 \pm (m+n-1)! \cdot y_m x^n \\ &= 1 \pm (m-1)! \cdot n! \cdot \tbinom{m+n-1}{n} y_m x^n. \end{aligned}$$

We have the following proposition whose proof follows from the above discussion.

**Proposition 3.5.** *The total Chern class of any element in the kernel of the realization map* $\rho : \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n) \to \widetilde{KO}(S^{2m} \wedge \mathbb{CP}^n)$ *is of the form*

(1) $1 + 2 \cdot (m-1)! \cdot \big[\sum_{i \geq 1} \binom{m+2i-1}{2i} (\sum_{k=1}^r b_k k^{2i}) y_m x^{2i}\big]$, *when $m$ is odd.*

(2) $1 - 2 \cdot (m-1)! \cdot \big[\sum_{i \geq 1} \binom{m+2i-2}{2i-1} (\sum_{k=1}^r b_k k^{2i-1}) y_m x^{2i-1}\big]$, *when $m, n$ are even.*

(3) $1 - (m-1)! \cdot \big[2\sum_{i=1}^{r+1} \binom{m+2i-2}{2i-1} (\sum_{k=1}^r b_k k^{2i-1}) y_m x^{2i-1} \pm \binom{m+n-1}{n} \cdot n! \cdot b_{r+1} y_m x^n\big]$, *when* $m \equiv 0 \pmod 4$ *and* $n \equiv 3 \pmod 4$, *or* $m \equiv 2 \pmod 4$ *and* $n \equiv 1 \pmod 4$.

(4) $1 - 2 \cdot (m-1)! \cdot \big[\sum_{i=1}^{r+1} \binom{m+2i-2}{2i-1} (\sum_{k=1}^r b_k k^{2i-1}) y_m x^{2i-1} \pm \binom{m+n-1}{n} \cdot n! \cdot b_{r+1} y_m x^n\big]$, *when* $m \equiv 0 \pmod 4$ *and* $n \equiv 1 \pmod 4$, *or* $m \equiv 2 \pmod 4$ *and* $n \equiv 3 \pmod 4$,

*for* $b_1, b_2, \cdots, b_r, b_{r+1} \in \mathbb{Z}$. ☐

**Remark 3.6.** In the above proposition, using that the fact that a binomial coefficient $\binom{s}{t}$ is even if $s$ is even and $t$ is odd, one can see that if $m$ is even and $n > 1$ then for any $a_1 \in \widetilde{K}(S^{2m} \wedge \mathbb{CP}^n)$ such that $\rho(a_1) = 0$ we have that $4 \cdot (m-1)!$ divides each Chern class $c_i(a_1)$.

**Proposition 3.7.** *If $S^{4p}\times\mathbb{CP}^n$ has a.c.s. then $2\cdot(2p-1)!$ divides $\chi(\mathbb{CP}^n)=(n+1)$.*

*Proof.* Suppose $a\in\widetilde{K}(S^{4p}\times\mathbb{CP}^n)$ gives an a.c.s. on $S^{4p}\times\mathbb{CP}^n$ with $a=a_1+a_2+a_3$. As we argued in the proof of Theorem 1.1, we have $a_2=0$. By Remark 3.6 we have that if $n>1$ then $4\cdot(2p-1)!$ divides each Chern class $c_i(a_1)$. Hence in the case $n>1$, we have that $4\cdot(2p-1)!$ divides the Euler characteristic $\chi(S^{4p}\times\mathbb{CP}^n)=2(n+1)$. In the case when $n=1$, we know that $S^{4p}\times\mathbb{CP}^1$ has a.c.s only when $p=1$. This completes the proof of the proposition. $\square$

Next we recall that for any element $a_3\in\widetilde{K}(\mathbb{CP}^n)$ such that $\rho(a_3)=[\tau_{\mathbb{CP}^n}]$, the total Chern class of $a_3$ has been described on p.130 of [13] as

$$c(a_3)=(1-x)^{n+1}(1\pm(n-1)!x^n)^{ud_{r+1}}\prod_{1\le k\le r}\Big(\frac{1+kx}{1-kx}\Big)^{d_k}, \tag{3}$$

where $d_i$'s are integers and

$$\begin{aligned} u &= 0 && \text{if } n \text{ is even,}\\ &= 1 && \text{if } n\equiv 3 \pmod 4,\\ &= 2 && \text{if } n\equiv 1 \pmod 4. \end{aligned}$$

We remark that there is a typographical error in the signs in the first two products of the right hand side of Equation 3 as expressed in [13]. For example one can easily compute $c(\eta^n)=(1+(n-1)!x^n)$ when $n=1,3$ whereas $c(\eta^2)=(1-x^2)$ when $n=2$.

We know that $S^{2m}\times\mathbb{CP}^1$ has a.c.s. if and only if $m=1,2$ and 3. Next we give the proof of Theorem 1.2.

**Proof of Theorem 1.2.** First note that $S^2\times\mathbb{CP}^n$, $S^6\times\mathbb{CP}^n$ have a.c.s. By Corollary 2.3 and Proposition 3.7, only thing that remains to complete the proof of the theorem is to show that for $q>0$ there is no a.c.s on $S^4\times\mathbb{CP}^{4q+1}$. Suppose $a\in\widetilde{K}(S^4\times\mathbb{CP}^{4q+1})$ gives an a.c.s. on $S^4\times\mathbb{CP}^{4q+1}$ with $a=a_1+a_2+a_3$. As observed in the proof of Theorem 1.1, we have $a_2=0$. The total Chern class of $a_1$ is given in (3) of Proposition 3.5 as

$$c(a_1)=1-2\sum_{i=1}^{2q+1}2i(\sum_{k=1}^{2q}b_kk^{2i-1})y_2x^{2i-1}\pm(4q+2)!\cdot b_{2q+1}y_2x^{4q+1}$$

where $b_i$'s are integers. The total Chern class of $a_3$ as described in Equation 3 is as follows

$$c(a_3)=(1-x)^{4q+2}(1\pm(4q)!x^{4q+1})^{2d_{2q+1}}\prod_{1\le k\le 2q}\Big(\frac{1+kx}{1-kx}\Big)^{d_k},$$

where $d_i$'s are integers. After simplifying we can write

$$c(a_3)=1+\sum_{i=1}^{2q}\binom{4q+2}{2i}x^{2i}+A$$

where $A$ is a polynomial in $x$ with even coefficients. As $c(a)=c(a_1)c(a_3)$, the top Chern class

$$c_{4q+3}(a)=(-2\sum_{i=1}^{2q+1}2i\binom{4q+2}{2i}\sum_{k=1}^{2q}b_kk^{2i-1}+h)y_2x^{4q+1} \tag{4}$$

where $h$ is a multiple of 8. We shall next prove that the coefficient

$$h_k:=-2\sum_{i=1}^{2q+1}2i\binom{4q+2}{2i}k^{2i-1},$$

of $b_k y_2 x^{4q+1}$ in Equation 4 is a multiple of 8. Clearly, when $k$ is even then $h_k$ is a multiple of 8. Next consider the case when $k$ is odd. Consider the following equality.

$$(4q+2)(1+k)^{4q+1} = \sum_{l=1}^{4q+2} \binom{4q+2}{l} l k^{l-1}.$$

As $k$ is odd, the left hand summation is a multiple of 4. The right hand summation can be decomposed into three parts as follows:

$$\textstyle\sum_{i=1}^{2q+1} 2i\binom{4q+2}{2i} k^{2i-1} \quad + \quad \sum_{j=1}^{q} \binom{4q+2}{2j-1}\Big[(2j-1)k^{2j-2} + (4q-2j+3)k^{4q-2j+2}\Big]$$
$$+ \quad \textstyle\binom{4q+2}{2q+1}(2q+1)k^{2q}.$$

The middle term in the above summation is a multiple of 4. To see that $\binom{4q+2}{2q+1}$ is a multiple of 4, we write

$$\binom{4q+2}{2q+1} = \frac{(4q+2)(4q+1)}{(2q+1)^2}\binom{4q}{2q}$$

and use the fact that $\binom{4q}{2q}$ is even which can be seen from the following equality

$$2^{4q} = (1+1)^{4q} = \sum_{i=0}^{4q} \binom{4q}{i}.$$

This completes the proof that each $h_k$ is a multiple of 8. Since $a$ gives an a.c.s. on $S^4 \times \mathbb{CP}^{4q+1}$, the top Chern class $c_{4q+3}(a)$ is multiple of 8. This implies that 8 divides the Euler characteristic $2(4q+2)$. But this is a contradiction. Hence there is no a.c.s. on $S^4 \times \mathbb{CP}^{4q+1}$. This completes the proof of the theorem. $\square$

The above construction helps us to give an explicit way to obtain almost complex structures on $S^{2m} \times \mathbb{CP}^n$ for few values of $m$ and $n$. In particular when $m = 2$ and $n = 3$ we show that $S^4 \times \mathbb{CP}^3$ has infinite number of almost complex structures. We fix an orientation on $S^{2m}$ and $\mathbb{CP}^n$ such that $e(S^{2m}) = -2y_m$ and $e(\mathbb{CP}^n) = (-1)^n(n+1)x^n$. We fix an orientation on $S^{2m} \times \mathbb{CP}^n$ arising by taking the orientation on each factor.

**Example 3.8.** Let $m = 1$ and $n = 1$. From Proposition 3.2 and 3.3, we get that $a = a_1 + a_2 + a_3 \in \widetilde{K}(S^2 \times \mathbb{CP}^1)$ gives an a.c.s. on $S^2 \times \mathbb{CP}^1 = S^2 \times S^2$ if and only if there are integers $d_1$, $d_2$ such that $a_1 = 0$, $a_2 = 2d_1\eta$ and $a_3 = [T_{\mathbb{CP}^1}] + 2d_2\eta$ and $c_2(a) = e(S^2 \times S^2) = (-2y_1)(-2x) = 4y_1x$. This gives the equation

$$4d_1(d_2 - 1) = 4,$$

which has the following two solutions: $d_1 = 1$, $d_2 = 2$ and $d_1 = -1$, $d_2 = 0$. Therefore we have the following two possibilities: $a = 2\eta + [T_{\mathbb{CP}^1}] + 4\eta$ and $a = -2\eta + [T_{\mathbb{CP}^1}]$. By stability property, we thus have that there are exactly two non-isomorphic a.c.s. on $S^2 \times \mathbb{CP}^1$. Hence the only two almost complex structures are $\bar{T}_{S^2} \oplus \bar{T}_{\mathbb{CP}^1}$ and $T_{S^2} \oplus T_{\mathbb{CP}^1}$. Here $\bar{T}_{S^2}$ is complex conjugate bundle of $T_{S^2}$ and $[\bar{T}_{\mathbb{CP}^1}] = [T_{\mathbb{CP}^1}] + 4\eta$. We note here that if we change the orientation on $S^2 \times S^2$, a similar argument will say that again it has exactly two almost complex structures which will be given by $T_{S^2} \oplus \bar{T}_{\mathbb{CP}^1}$ and $\bar{T}_{S^2} \oplus T_{\mathbb{CP}^1}$. The existence of exactly two almost complex structures on $S^2 \times S^2$ was also observed by Sutherland in [9].

**Example 3.9.** Let $m = 1$ and $n = 2$. As in Example 3.8, an element $a \in \widetilde{K}(S^2 \times \mathbb{CP}^2)$ gives an a.c.s. on $S^2 \times \mathbb{CP}^2$ if and only if there are integers $b_1$, $d_1$ and $d_2$ such that $a_1 = b_1(g(H-1) - \overline{g(H-1)})$,

$a_2 = 2d_1\eta$, $a_3 = [T_{\mathbb{CP}^2}] + d_2(H - \bar{H})$ and $c_3(a) = -6y_1x^2$. This gives the following equations

$$b_1 + d_1(-4d_2 + 4\tbinom{d_2}{2} + 3) = -3 \qquad \text{when } d_2 \geq 0$$

$$b_1 + d_1(-8d_2 + 4\tbinom{-d_2}{2} + 3) = -3 \qquad \text{when } d_2 < 0.$$

Clearly, we shall get infinite number of solutions of the above equations and by stability property, we have infinite number of almost complex structures on $S^2 \times \mathbb{CP}^2$. The solution $b_1 = 0$, $d_1 = -1$ and $d_2 = 0$ corresponds to the a.c.s. given by the holomorphic tangent bundle $T_{S^2} \oplus T_{\mathbb{CP}^2}$. Similarly, if we change the orientation on $S^2 \times \mathbb{CP}^2$, we shall again get infinite number of almost complex structures.

**Example 3.10.** Let $m = 2$ and $n = 3$. Going along the same line of arguments as we did in last two examples we observe that number of almost complex structures on $S^4 \times \mathbb{CP}^3$ is in one to one correspondence with the number of solutions of the following equation

$$b_1(4 - 3d_1 + 2\tbinom{d_1}{2}) + 3!b_2 = -1 \qquad \text{when } d_1 \geq 0$$

$$b_1(4 - 5d_1 + 2\tbinom{-d_1}{2}) + 3!b_2 = -1 \qquad \text{when } d_1 < 0.$$

For each $k \in \mathbb{Z}$, by taking $b_1 = -7 + 6k$, $b_2 = 1 - k$ and $d_1 = 1$ we get infinite number of solutions of the above equation and thus this gives infinite number of almost complex structures on $S^4 \times \mathbb{CP}^3$. Again the reverse orientation on $S^4 \times \mathbb{CP}^3$ has infinite number of almost complex structures. We remark that the existence of an a.c.s. on $S^4 \times \mathbb{CP}^3$ also follows from Theorem 2 of [6] and this was observed by Tang in [11].

We end this section with a result on nonexistence of a.c.s. on the Dold manifold $D(m, n)$. It is known (see [3]) that $D(m, n)$ is orientable if and only if $m + n$ is odd. So an even dimensional orientable Dold manifold is of the form $D(2p, 2q + 1)$ for some $p, q \geq 0$. By considering the 2-fold covering map $S^{2p} \times \mathbb{CP}^{2q+1} \to D(2p, 2q + 1)$, we note that the nonexistence of an a.c.s. on $S^{2p} \times \mathbb{CP}^{2q+1}$ will imply the nonexistence of an a.c.s. on $D(2p, 2q+1)$. We have the following result.

**Theorem 3.11.** *Let $p > 0$. Then $D(2p, 2q + 1)$ has no a.c.s. if one of the following is true*

1. *$p$ is odd.*
2. *$p \equiv 0 \pmod 4$ and $(q + 1)$ is not a multiple of $2^{r-2} \cdot (p - 1)!$, where $2^r$ is the maximum power of 2 dividing $p$.*
3. *$p \equiv 2 \pmod 4$ and $(q + 1)$ is not a multiple of $(p - 1)!$.*
4. *$p = 2$ and $q$ is even.*

*Proof.* Statement (1) follows from Theorem 3.2 of [12]. The proof of statements (2), (3) and (4) follow from Theorem 1.1, Proposition 3.7 and Theorem 1.2 respectively. □

From the above theorem it is clear that $D(2p, 2q + 1)$ does not admit an a.c.s. if $p$ is odd or $q$ is even.

**Acknowledgement:** We are grateful to Aniruddha Naolekar for several valuable discussions and useful comments.

## References

[1] Borel, A., Serre, J. P., *Groupes de Lie et puissances réduites de Steenrod*, Amer. J. Math. **75** (1953), 409-448.
[2] Datta, B., Subramanian, S., *Nonexistence of almost complex structures on products of even-dimensional spheres*, Topology Appl. **36** (1990), no. 1, 39-42.
[3] Dold, A., *Erzeugende der Thomschen Algebra $\mathfrak{N}$*, Math. Z. **65** (1956), 2535.
[4] Fujii, M., *KU-groups of Dold manifolds*, Osaka J. Math. **3** (1966), 49-64.

[5] Fujii, M., *KO-groups of projective spaces*, Osaka J. Math. **4** (1967), 141-149.

[6] Heaps, T., *Almost complex structures on eight- and ten-dimensional manifolds*, Topology **9** (1970), 111-119.

[7] Husemoller, D., *Fibre bundles* Third edition. Graduate Texts in Mathematics, 20. Springer-Verlag, New York, 1994.

[8] Kobayashi, T., *Note on almost complex structures on products of lens spaces* Math. Proc. Cambridge Philos. Soc. **96** (1984), no. 1, 81-83.

[9] Sutherland, W. A., *A note on almost complex and weakly complex structures*, J. London Math. Soc. **40** (1965), 705-712.

[10] Tanaka, R., *On trivialities of Stiefel-Whitney classes of vector bundles over iterated suspension spaces*, Homology, Homotopy Appl. **12** (2010), no. 1, 357-366.

[11] Tang, Z., *Almost complex structures on* $S^{2m} \times \mathbb{CP}^2$ *and* $S^{2m} \times \mathbb{CP}^3$, Chinese Sci. Bull., **37** (1992), no. 24, 2025-2028.

[12] Tang, Z., *Nonexistence of almost complex structures on Dold manifolds*, Sci. China Ser. A **39** (1996), no. 9, 919-924.

[13] Thomas, A., *Almost complex structures on complex projective spaces*, Trans. Amer. Math. Soc. **193** (1974), 123-132.

[14] Thomas, A., *Counting almost complex structures on products of lens spaces*, J. London Math. Soc. (2) **7** (1974), 761-768.

Stat-Math Unit, Indian Statistical Institute, 8th Mile, Mysore Road, RVCE Post, Bangalore 560059, INDIA.

*E-mail address*: `chakraborty.prateep@gmail.com`

Stat-Math Unit, Indian Statistical Institute, 8th Mile, Mysore Road, RVCE Post, Bangalore 560059, INDIA.

*E-mail address*: `thakur@isibang.ac.in`